

Controllable Josephson-Like Tunneling in Two-Component Bose-Einstein Condensates Coupled with Microwave via Feshbach Resonance and Trapping Potential

Bo Xiong^{1,2}, Weiping Zhang², W. M. Liu¹

¹*Beijing National Laboratory for Condensed Matter Physics,*

Institute of Physics, Chinese Academy of Sciences, Beijing 100080, China and

²*Key Laboratory of Optical and Magnetic Resonance Spectroscopy (Ministry of Education),
Department of Physics, East China Normal University, Shanghai 200062, China*

(Dated: February 6, 2008)

We put forward a scheme for controlling Josephson-like tunneling in two-component Bose-Einstein condensates coupled with microwave field via Feshbach resonance and tuning aspect ratio of trapping potential. We prove how to realize a perfect periodic oscillation from a fast damped and irregular oscillation on relative number of atoms in future experiment. In particular, intensity of Josephson-like tunneling can be successfully controlled through controlling speed of recovering the initial value of intra-atomic interaction and aspect ratio of trapping potential. Interestingly, we find that relative number of atoms represents two different types of oscillation in respond to periodic modulation of attractive intra-atomic interaction.

PACS numbers: 03.75.-b, 67.40.-w, 39.25.+k

The existence of a Josephson current is a direct manifestation of macroscopic quantum phase coherence [1] and have numerous important applications in condensed matter physics, quantum optics and cold atom physics, for example, precision measurement, quantum computation. The physical origin of the Josephson current is the temporal interference of the two systems which must both have a well defined quantum phase and a different average energy per particle, respectively. Recently, the experimental realization of multi-component Bose-Einstein condensates (BECs) of weakly interacting alkali atoms has provided a route to study Josephson effect [2, 3, 4, 5, 6, 7, 8, 9, 10] in a controlled and tunable way by means of Feshbach resonance [11, 12, 13, 14, 15, 16] so far unattainable in charged systems. The physical origin of the low-energy Feshbach resonances is the low-energy binary collisions described by the difference of the initial and intermediate state energies which can be effectively altered through variations of the strength of an external magnetic field.

In this Letter, we put forward a scheme for controlling Josephson-like tunneling in two-component BECs coupled with microwave field via Feshbach resonance and tuning aspect ratio of trapping potential. we find that, through time-dependent tuning attractive intra-atomic interaction, a perfectly periodic oscillation can be obtained successfully. Especially, by controlling speed of recovering the initial value of attractive intra-atomic interaction and aspect ratio of trapping potential, we can successfully control intensity of Josephson-like tunneling. Furthermore, relative number of atoms represents two different types of oscillation in respond to periodic modulation of attractive intra-atomic interaction.

We consider a system of longitudinal elongated two-component BECs coupled by the microwave field with the effective Rabi frequency Ω and finite detune δ , where the Ω and δ are independent upon time and space coordi-

nate as experimental case [3]. At zero temperature, this system can be described by macroscopic wave function ψ_j (normalized to unity, i.e., $\int (|\psi_1|^2 + |\psi_2|^2) dz = 1$) which obey the dimensionless one-dimensional Gross-Pitaevskii equations (GPEs) as [6, 7]

$$i \frac{\partial}{\partial t} \psi_j(z, t) = \left[-\frac{\partial^2}{\partial z^2} + V(z) \right] \psi_j(z, t) + [G_{jj}(t)|\psi_j|^2 + G_{jk}(t)|\psi_k|^2] \psi_j(z, t) + (-1)^{j+1} \frac{\delta}{\omega_{\perp}} \psi_j(z, t) + \frac{\Omega}{\omega_{\perp}} \psi_k(z, t), \quad (1)$$

where $j, k = 1, 2$, $j \neq k$, and $\omega_{\perp}$ is radial harmonic trapping frequency, time and coordinate unit are $2/\omega_{\perp}$ and $a_{\perp} = \sqrt{\hbar/m\omega_{\perp}}$, respectively, and $V(z)$ is the longitudinal confining potential, $G_{jj} = (4N\hbar a_{jj})/(ma_{\perp}^3\omega_{\perp})$ and $G_{jk} = (4N\hbar a_{jk})/(ma_{\perp}^3\omega_{\perp})$ are dimensionless intra- and inter-species effective atomic interaction potentials, respectively, where a_{jj} and a_{jk} are corresponding s-wave scattering lengths, respectively, N is the total atomic number, and m is the atomic mass assumed the same for both two species.

In the following discussion, we consider two-hyperfine spin states of ^7Li bose atom which the magnitude and sign of the atomic interaction can be tuned to any value via a magnetic-field dependent Feshbach resonance. And we choose dimensionless $V(z) = \gamma^2 z^2$, where $\gamma = \omega_z/\omega_{\perp}$ is aspect ratio of trapping potential, ω_z is longitudinal harmonic trapping frequency and $\omega_{\perp} = 2\pi \times 625$ Hz (so, the time and coordinate units are 0.51 ms and $a_{\perp} = 1.51$ μm respectively). Moreover, considering the condensates will be unstable and collapse when the number of particles is large enough [17], so we choose $N = 6000$, which provides a safe range of parameters for avoiding instability and collapse occurring. And the effective Rabi frequency and finite detune are chosen as $\Omega = 1.5\omega_{\perp}$ and $\delta = 0.5\omega_{\perp}$ considering experimental feasibility.




FIG. 1: (Color online) Temporal evolution of atomic density and relative number of atoms N_d for initial Gaussian wave function in response to various aspect ratio of trapping potential γ . (a) Corresponding to attractive atomic interactions for the parameters as $G_{11} = G_{22} = 4G_{12} = -10$. (b) Corresponding to repulsive atomic interactions for the parameters as $G_{11} = G_{22} = 4G_{12} = 10$.

First of all, we consider the simplest case, i.e., intra- and inter-species effective atomic interaction potentials are time-independent attractive or repulsive case, which can be easily realized using Feshbach resonance experimentally. Here, we have chosen $a_{11} = a_{22}$ is just considering the experimental feasibility and assumed that tuning atomic interaction wouldn't destroy the condensates in this system. We denote relative number of atoms between two species as N_d , i.e., $N_d = N_1 - N_2$, where $N_1 = \int |\psi_1|^2 dz$ and $N_2 = \int |\psi_2|^2 dz$, respectively. We study Josephson-like tunneling in this case and the results are shown in Fig. 1, which is based on numerical simulation of Eqs. (1) for the initial Gaussian wave function under the various aspect ratio of trapping potential. As shown in Fig. 1a corresponding to attractive atomic interactions, during the evolution process, exchange of atoms is always damped very fast and almost suppressed completely irrespective of aspect ratio of trapping potential. But, on the contrary, in repulsive atomic interactions corresponding to Fig. 1b, exchange of atoms vary significantly in response to various aspect ratio of trap-




FIG. 2: (Color online) Temporal evolution of N_d for initial Gaussian wave function in response to various aspect ratio of trapping potential γ , which is based on numerical simulation of Eqs. (3) for the case $\Sigma(\eta) = 0$. (a) Corresponding to attractive atomic interactions for the parameters as $G_{11} = G_{22} = 4G_{12} = -10$. (b) Corresponding to repulsive atomic interactions for the parameters as $G_{11} = G_{22} = 4G_{12} = 10$.

ping potential. Detailed analyses of above phenomena will be given in the following.

From Eqs. (1) and taking boundary condition into account, it is not difficult to derive following equations,

$$\frac{dN_d}{dt} = \frac{4\Omega}{\omega_{\perp}} \int (\sqrt{\rho_1 \rho_2} \sin \phi_d) dz, \quad (2)$$

where wave functions have been chosen as $\psi_1 = \sqrt{\rho_1} \exp(i\phi_1)$, $\psi_2 = \sqrt{\rho_2} \exp(i\phi_2)$, respectively, and $\phi_d = \phi_1 - \phi_2$. From Fig. 1a, we can see that maximum ρ_1 will be much smaller than maximum ρ_2 very quickly for all aspect ratio of trapping potential, so maximum $\sqrt{\rho_1 \rho_2}$ will decrease rapidly. Moreover, if considering well localized density profiles of two species as shown in Fig. 1a, from Eqs. (2), we can immediately conclude that amplitude of oscillation of N_d will be damped very fast and suppressed completely in this case. But from Fig. 1b corresponding to repulsive atomic interactions, we can see that the atomic densities of two species vary significantly in response to various aspect ratio of trapping potential and represent nonlocalized profiles, so time evolution of amplitudes of N_d represents very different behavior from attractive case.

As pointed out above, N_d will represent a very irregular and fast damped oscillation in time-independent attractive atomic interactions and represent a more irregular oscillation in time-independent repulsive case. But, from the experimental point of view, valuable and applicable systems usually have regular and stable characteristics rather than irregular and unstable one. Fortunately, we




FIG. 3: (Color online) Temporal evolution of relative number of atoms N_d for initial Gaussian wave function in response to various aspect ratio of trapping potential γ . (a) Corresponding to attractive atomic interactions for the parameters as $G_{12} = -2.5$, $\tau = 105$, $t_1 = 0.5t_2 = 280$. (b) Corresponding to repulsive atomic interactions for the parameters as $G_{12} = 2.5$, $\tau = 105$, $t_1 = 0.5t_2 = 280$.

find out that, through time-dependent tuning atomic interaction via Feshbach resonance, regular and stable oscillation of N_d can be successfully realized in the attractive case, but not be realized in the repulsive case. Now, let us to understand physics from both theoretical analysis and numerical simulation aspects.

In order to study how to control Josephson-like tunneling in this system, we diagonalize linear parts of Eqs. (1) by means of applying a unitary transformation U on both sides of Eqs. (1), then it is transformed into following form,

$$i \frac{\partial}{\partial t} \varphi_{\pm}(z, t) = \left[-\frac{\partial^2}{\partial z^2} + V(z) \right] \varphi_{\pm}(z, t) + \zeta_1^{\pm} |\varphi_{\pm}|^2 \varphi_{\pm}(z, t) + \zeta_2^{\pm} |\varphi_{\mp}|^2 \varphi_{\pm}(z, t) + \Sigma(\eta), \quad (3)$$

where

$$U \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = \begin{pmatrix} \varphi_+ e^{i\eta t} \\ \varphi_- e^{-i\eta t} \end{pmatrix} \equiv \begin{pmatrix} \Psi_+ \\ \Psi_- \end{pmatrix}, U = \begin{pmatrix} \sin \theta & -\cos \theta \\ \cos \theta & \sin \theta \end{pmatrix},$$

$$\Sigma(\eta) = \zeta_3^{\pm} \varphi_{\pm}^2 \varphi_{\mp}^* e^{\pm 2i\eta t} + \zeta_4^{\pm} |\varphi_{\pm}|^2 \varphi_{\mp} e^{\mp 2i\eta t} + \zeta_5^{\pm} \varphi_{\mp}^2 \varphi_{\pm}^* e^{\mp 4i\eta t} + \zeta_6^{\pm} |\varphi_{\mp}|^2 \varphi_{\pm} e^{\mp 2i\eta t},$$

here $\zeta_1^{\pm} = [(3G_{11} + 2G_{12} + 3G_{22}) \mp 4(G_{11} - G_{22}) \cos 2\theta + (G_{11} - 2G_{12} + G_{22}) \cos 4\theta]/8$, $\zeta_2^{\pm} = [(G_{11} + 2G_{12} + G_{22}) - (G_{11} - 2G_{12} + G_{22}) \cos 4\theta]/4$, $\zeta_3^{\pm} = [2(G_{11} - G_{22}) \sin 2\theta \mp (G_{11} - 2G_{12} + G_{22}) \sin 4\theta]/8$, $\zeta_4^{\pm} = [2(G_{11} -$




FIG. 4: (Color online) Temporal evolution of N_d for initial Gaussian wave function in response to (a) various speed of recovering the initial value of intra-atomic interaction for the parameters as $\gamma = 0$, $G_{12} = -2.5$, $\tau = 21$, $t_1 = 56$, and (b) various aspect ratio of trapping potential for the parameters as $G_{12} = -2.5$, $\tau = 21$, $t_1 = 0.83t_2 = 56$.

$G_{22}) \sin 2\theta \mp (G_{11} - 2G_{12} + G_{22}) \sin 4\theta]/4$, $\zeta_5^{\pm} = [(G_{11} - 2G_{12} + G_{22}) \sin^2 2\theta]/4$, $\zeta_6^{\pm} = [2(G_{11} - G_{22}) \sin 2\theta \pm (G_{11} - 2G_{12} + G_{22}) \sin 4\theta]/8$, and $\theta = (1/2) \tan^{-1}(\Omega/\delta)$, $\eta = [(\delta/\omega_{\perp})^2 + (\Omega/\omega_{\perp})^2]^{1/2}$. Now, we can see that, comparing with the original Eqs. (1), we introduce a new term denoted by $\Sigma(\eta)$ which is exponential time dependence, and it is not included in regular coupling nonlinear Schrödinger equations. It must be pointed out that $\Sigma(\eta)$ depends not only on the effective atomic interaction potentials G_{jk} ($j, k = 1, 2$), but on the effective Rabi frequency Ω and finite detune δ , i.e., θ and η .

From the expression of $\Sigma(\eta)$, we can find that, if $\varphi_{\pm}(z, t)$ vary slowly over time period π/η , then the effects of $\Sigma(\eta)$ on Josephson-like tunneling can be averaged out over time period π/η during the evolution process. But if $\varphi_{\pm}(z, t)$ vary significantly over time period π/η , then $\Sigma(\eta)$ will strongly affect Josephson-like tunneling. In the following, we will investigate effects of $\Sigma(\eta)$ on Josephson-like tunneling in detail.

Firstly, we consider the special case that $\Sigma(\eta) = 0$. From Eqs. (3) and transformation relation between original and transformed wave function, we can obtain that N_d will represent a large-amplitude periodic oscillation in both attractive and repulsive atomic interactions. Above conclusions are well confirmed by Fig. 2, which is based on numerical simulation of Eqs. (3) and then applying inverse unitary transformation U^{-1} on $\Psi_{\pm}$ in order to obtain original wave function. So, it is reasonable to infer that the reasons for fast damped and irregular oscillation of N_d are mainly due to the effects of $\Sigma(\eta)$ on Josephson-like tunneling.

But, key question is how to efficiently suppress effects of $\Sigma(\eta)$ on Josephson-like tunneling in order to obtain

perfectly periodic oscillation of N_d . In the following, we prove that, only in the attractive case, through time-dependent tuning intra-atomic interaction via Feshbach resonance, we can successfully control the effects of $\Sigma(\eta)$ and obtain a perfectly periodic oscillation on relative number of atoms. Considering the experimental feasibility, in the following case, we choose time-dependent intra-atomic interaction as [18, 19, 20]

$$G_{jj}(t) = \begin{cases} G_{12} \exp(t/\tau), & 0 \leq t \leq t_1; \\ G_{12} \exp[(t_2 - t)/\tau], & t_1 < t \leq t_2; \\ G_{12}, & t > t_2. \end{cases} \quad (4)$$

where $j = (1, 2)$, $G_{11}(t) = G_{22}(t)$ and G_{12} is time-independent parameter determining inter-atomic interaction, τ determines speed of tuning intra-atomic interaction depending on the values of t_1 and t_2 at fixed minimum $G_{jj}(t)$. The results are shown in Fig. 3. It is very interesting that N_d finally represents a perfectly periodic oscillation in the attractive case, but not in the repulsive case. Above phenomena are mainly due to successfully suppress the effects of $\Sigma(\eta)$ on Josephson-like tunneling in the attractive case, but not in the repulsive case, which underlying physics is just because $\varphi_{\pm}(z, t)$ vary significantly over time period π/η in the repulsive case. So, in the following, we will concentrate on attractive case. It is important to point out that, in the attractive case, the amplitude of N_d is strongly dependent upon aspect ratio of trapping potential as shown in Fig. 3a. Based upon the above analysis, we put forward a scheme for controlling the amplitude of oscillation through controlling speed of recovering the initial value of intra-atomic interaction (i.e., changing value of t_2 but fixing other parameters in Eqs. (4)) and tuning aspect ratio of trapping potential.

The results are shown in Fig. 4. It is very interesting to note that oscillatory amplitude of N_d can be controlled successfully not only via Feshbach resonance but tuning aspect ratio of trapping potential. Above phenomena can be well understood as follows: If $\varphi_{\pm}(z, t)$ vary slowly over time period π/η , then the effects of $\Sigma(\eta)$ on Josephson-like tunneling can be effectively suppressed, and N_d will represent a perfectly periodic oscillation which amplitude is determined by atomic density of two species.

We also study Josephson-like tunneling in response to periodic modulation of attractive interaction [12] and various aspect ratio of trapping potential. In this case, attractive atomic interactions have been chosen as $G_{11}(t) = G_{22}(t) = G_{12}[1 + 8\sin^2(0.01t)]$, where $G_{12} = -2.5$. As shown in Fig. 5, it is very interesting to note that N_d represents two different types of oscillation, which one represents small frequency behavior originated from periodic modulation of attractive interaction, but the other represents large frequency behavior originated from coupling microwave. Above effects provide a promising protocol to realize a new type of switch




FIG. 5: (Color online) Temporal evolution of relative number of atoms N_d for initial Gaussian wave function in response to periodic modulation of attractive intra-atomic interaction and various aspect ratio of trapping potential for the parameter as $G_{12} = -2.5$.

controlled by periodically managed Feshbach resonance in future experiments.

In conclusion, we prove how to successfully control Josephson-like tunneling in attractive two-component BECs coupled by microwave field via Feshbach resonance in respond to various aspect ratio of trapping potential. Recent developments of controlling the scattering length and realization of multi-component BECs in the experiments allow for the experimental investigation of our prediction in future.

This work was supported by the NSF of China under 10474055, 90406017, 10588402, 60525417; the NKBRF of China under 2005CB724508; the STCM of Shanghai under 05PJ14038, 06JC14026, 04DZ14009.

-
- [1] A. Barone, G. Paterno, Physics and Applications of the Josephson Effect (Wiley, New York, 1982).
 - [2] D. S. Hall et al., Phys. Rev. Lett. **81**, 1539 (1998); ibid **81**, 1543 (1998).
 - [3] M. R. Matthews et al., Phys. Rev. Lett. **81**, 243 (1998); ibid **83**, 3358 (1999).
 - [4] A. J. Leggett, Rev. Mod. Phys. **73**, 307 (2001).
 - [5] B. P. Anderson et al., Science **282**, 1686 (1998).
 - [6] J. Williams et al., Phys. Rev. A **59**, R31 (1999).
 - [7] Q. H. Park et al., Phys. Rev. Lett. **85**, 4195 (2000).
 - [8] F. S. Cataliotti et al., Science **293**, 843 (2001).
 - [9] S. Kohler et al., Phys. Rev. Lett. **89**, 060403 (2002).
 - [10] Y. Shin et al., Phys. Rev. Lett. **95**, 170402 (2005).
 - [11] H. Saito et al., Phys. Rev. Lett. **90**, 040403 (2003).
 - [12] P. G. Kevrekidis et al., Phys. Rev. Lett. **90**, 230401 (2003).
 - [13] D. E. Pelinovsky et al., Phys. Rev. Lett. **91**, 240201 (2003).
 - [14] Víctor M. Pérez-García et al., Phys. Rev. Lett. **92**, 220403 (2004).
 - [15] V. V. Konotop et al., Phys. Rev. Lett. **94**, 240405 (2004).
 - [16] M. Matuszewski et al., Phys. Rev. Lett. **95**, 050403 (2005).
 - [17] A. E. Muryshev et al., Phys. Rev. Lett. **60**, R2665 (1999).
 - [18] S. Inouye et al., Nature **392**, 151 (1998).
 - [19] J. L. Roberts et al., Phys. Rev. Lett. **85**, 728 (2000).

- [20] K. E. Strecker, et al., *Nature* **417**, 150 (2002).